

Effect of Priming and Storage on Seed Quality of Chilli

P. Divya and S. NirmalaDevi

Department of Olericulture,

Kerala Agricultural University, Vellanikkara, Thrissur.

E mail: divya3233@gmail.com

Abstract

The present research was undertaken to study the effect of priming and storage of seeds of chilli freshly harvested seeds of chilli var. Anugraha were stored under ambient condition for 12 months. Seed samples were drawn after 3, 6 and 9 months of storage and subjected to priming treatments *viz.* Control (P1), Water soaking (P2), NaCl (10- 5M) (P3), CaCl₂ (10-5M), (P4 5), PEG 6000 (-1.5MPa) (P6) and (10g/kg seed) (P7). The primed seeds were stored in two types of storage containers i.e. cloth bag (S1) and polythene bag (700 gauge) (S2). The seeds primed with PEG 6000 (-1.5MPa) and stored in polythene bag (700 gauge) (S2) retained high hand, the seed quality parameters of control seeds stored in cloth bag was declined.

Key words: *chilli seed, germination, priming, storage containers*

Introduction

Chilli (*Capsicum annuum*) is one of the important vegetable crops of the world and is widely cultivated throughout the tropical and subtropical countries. There has been a great competition in the world market for chilli and hence it is necessary to increase production and productivity. The availability of quality seed is essential to achieve optimum crop stand in the field. The most important aspect in the seed production programme is maintenance of regular supply of high quality seeds to farmers. Under ambient conditions of storage, chilli loses its viability within a year.

Strategies for improving the growth and development of crop species have been investigated for many years. In field crop seed priming techniques have been used to improved germination characteristics. During priming, seeds are partially hydrated so that pre-germination metabolic activities proceed, while radicle protrusion is prevented, then are dried back to the primary moisture level (McDonald, 1999). Various priming treatments have been employed to increase the rate and uniformity of seed germination (Bradford, 1986). Common priming techniques include osmo priming (soaking seeds in osmotic solutions such as polyethylene glycol), halo priming (soaking seeds in salt solutions), hydro priming (soaking seeds in water), hormone priming (soaking

seeds in hormone solutions) and other various chemical solutions. Priming contributes to significant improvement in seed germination and seedling growth in different plant species.

The most important aspect to narrow down the gap between potential and farm level yield is the use of quality seed as it ensures better germination and there by better yield. Although seed quality is governed by genetic make-up, but commonly the quality of seeds is deteriorated during storage. The performance of the primed seeds is mainly affected by the storage conditions. Consequently, this research studied the possibilities to identify priming and storage procedures of primed chilli seeds with the primary objective of maintaining the beneficial effect of this treatment.

Materials and methods

A freshly harvested seed lot of chilli (*Capsicum annuum* L.) cultivar Anugraha with moisture content of 8 % (on dry weight basis) was obtained from Department of Olericulture, KAU, Vellanikkara.

Priming treatment: seeds were surface sterilized by dipping in sodium hypochlorite(5%) solution for five minutes and dried on filter paper .The surface sterilized seeds subjected to various priming treatments *viz.* Water soaking (P2), NaCl (10-5M) (P3), CaCl₂ (10-5M), (P4

5), KNO₃ (150 ppm) (P5) PEG 6000 (-1.5 MPa) (P6) and *Pseudomonas fluorescens* (10g/kg seed) (P7). After respective priming treatment for 3 hours seeds were washed with distilled water and dried at room temperature on filter paper in shade. Primed seeds were then packed in cloth bag and polythene bag (700 gauge) and stored under ambient condition.

$$\text{Rate of germination} = \frac{X_1(X_2 - X_1) X_n - (X_n - 1)}{Y_1 Y_2 Y_n}$$

Where,

X_n – Number of seeds germinated at nth count

Y_n – Number of days from sowing to nth count

Electrical conductivity (dsm⁻¹)

Five grams of seeds were taken at random from each treatment and were soaked in HgCl₂ 0.10% solution for a minute and washed with distilled water for five times. Then the seeds were soaked in 25.00 ml of distilled water for 24 hr at room temperature. The seed leachate was collected by decanting and the volume was made up to 25.00 ml by adding distilled water and the electrical conductivity of seed leachate was measured in an electrical conductivity bridge (ELICO) with cell constant of 1.00. The estimations were done in four replications and mean value was expressed in dsm⁻¹ (Presley, 1958).

Vigour indices

The vigour index (V I) was calculated by adopting the method suggested by Abdul-Baki and Anderson (1973) and was expressed as pure number. Vigour index (V I) I = (germination (%) x seedling dry weight) vigour index (V I) II = germination (%) x (seedling length in cm).

Results and discussion

The seeds subjected to priming, after 3 months of storage, with PEG 6000 (1.5 MPa) and stored in polythene bag (700 gauge) recorded the maximum germination (56.3 %), speed of germination (8.2), vigour indices and minimum electrical conductivity of the seed leachate (0.704 dsm⁻¹). It may be due to the fact that during osmopriming the transition of seeds from dry stage to germination represses the antioxidant pathways that involve CAT (catalase) and SOD (superoxide dismutase) enzymes and stimulates another pathway involving ascorbate peroxidase (APX) required for germination and seedling establishment. In control where no priming was done germination (30.54%), speed of germination (4.07), and vigour indices were low which may be due to excessive leakage of chemicals from the seeds. This is evident from the high value for electrical conductivity in control.

The results are in conformity with the findings of Delouche (1973) who has reported that the increase in electrical conductivity in unprimed seeds was due to rapid loss of electrolytes from the seed due to membrane damage during imbibitions. After 6 months of storage, when the seeds were treated with PEG 6000 (-1.5 MPa) and stored in polythene bag (700 gauge) the maximum germination (55.3 %), speed of germination (7.5), vigour indices and minimum electrical conductivity of the seed leachates (0.443 dsm⁻¹) were recorded. The electrical conductivity was found to be high in seeds stored in cloth bag (S1) when compared to that stored in polythene bag (S2). This may be due to alterations in membrane integrity that led to enhanced exudation of leachates. Similar results were reported in soybean seeds by (Schoettle and Leopold, 1984). Doijode (1986) and Karivaratharaju and Palanisamy (1987) also observed increase in leaching of electrolytes in Okra and tomato seed respectively with increased period of storage.

Table 1: Effect of priming on quality of Chilli seeds primed at 3, 6, 9 months after storage

Treatments	Germination (%)			Speed of germination			Electrical conductivity			Vigour index 1			Vigour index 2		
	3 M	6 M	9 M	3 M	6 M	9 M	3 M	6 M	9 M	3 M	6 M	9 M	3 M	6 M	9 M
P ₁ S ₁	30.54	9.33	0.33	4.07	1.36	2.00	0.78	0.92	0.95	426.0	10807	21.00	315.8	92.71	19.20
P ₁ S ₂	38.25	15.33	1.19	5.12	2.29	6.67	0.73	0.86	0.93	598.7	189.80	73.33	426.2	160.97	66.00
P ₂ S ₁	45.92	38.33	3.66	6.27	5.21	26.33	0.72	0.54	0.83	746.7	558.13	357.00	537.5	489.87	301.23
P ₂ S ₂	50.00	45.00	4.31	6.78	6.04	31.33	0.69	0.53	0.82	824.3	679.33	424.00	604.7	592.38	373.08
P ₃ S ₁	47.92	46.20	4.16	6.73	6.05	30.00	0.65	0.52	0.82	738.2	694.00	410.50	595.1	582.72	348.10
P ₃ S ₂	52.13	5.93	4.62	7.30	6.89	32.83	0.62	0.51	0.80	821.9	814.27	454.67	673.8	672.67	397.43
P ₄ S ₁	45.46	42.73	4.33	6.25	5.60	31.33	0.69	0.49	0.80	682.3	662.73	424.17	567.7	499.89	417.38
P ₄ S ₂	49.92	46.87	4.90	6.96	6.37	34.67	0.66	0.47	0.79	783.0	738.40	4855.33	644.3	593.58	471.65
P ₅ S ₁	48.92	40.73	4.50	6.72	5.76	31.67	0.73	0.47	0.81	733.2	640.00	450.33	607.8	531.47	417.75
P ₅ S ₂	52.50	45.27	5.09	7.14	6.48	35.83	0.69	0.46	0.80	834.9	737.07	531.67	672.8	601.27	484.98
P ₆ S ₁	54.46	50.47	5.32	7.82	7.18	34.33	0.64	0.46	0.79	862.3	806.93	512.67	718.2	662.71	495.80
P ₆ S ₂	58.17	55.33	5.79	8.23	7.52	41.00	0.59	0.44	0.79	962.3	933.80	636.67	824.5	749.88	572.08
P ₇ S ₁	38.25	23.40	1.53	5.24	3.38	10.50	0.73	0.55	0.91	671.2	393.40	117.67	553.2	324.80	115.20
P ₇ S ₂	43.13	29.93	1.53	5.68	3.89	11.50	0.71	0.54	0.90	776.3	521.80	126.50	644.8	425.79	129.27

There was significant difference between treatments for percent germination (%), speed of germination and vigour indices when priming was done 9 months after storage. Similar result was also reported by Venkatasubramanian and Umarani (2010). Seeds treated with PEG 6000 (-1.5 MPa) and stored in polythene bag recorded highest percent germination (37.6 %), speed of germination (5.5) The low germination percent of unprimed seeds is due to high moisture content of seeds and high electrical conductivity of the seed leachate which gives an indication of poor quality of the seeds. High germination in seeds primed with PEG 6000 (-1.5 MPa) can be expected because of the low moisture content and low electrical conductivity of the seeds. Even though the germination (%) for the best treatment was less than 60 % (minimum standard for certified seeds) this will be useful in getting the precious seeds of the germplasm germinated even after 12 months of storage under ambient conditions.

The seed quality parameters of stored seeds (3, 6 and 9 months) recorded higher values of percent germination

and seedling vigour indices immediately after priming with PEG 6000 (-1.5 MPa). The unprimed (control) seeds maintained minimum seed certification standards for certified seeds only up to sixth month. The seed primed with PEG 6000 (-1.5 M Pa) and stored in polythene bag (700 gauge) after 3 months and 6 months of storage (ambient conditions) maintained seed certification standards up to eighth month and ninth month of storage respectively.

Conclusion

Loss of seed vigour and viability was said to be associated with the ageing phenomenon and results in poor stand and performance in the field which can be managed by priming and improving on storage conditions for promoting vigour, viability, storability and performance in chilli. Priming treatments exhibited marked influence on seed germination and seedling growth of chilli. The seeds primed with PEG 6000 (-1.5MPa) and stored in polythene bag (700 gauge) (S2) retained high hand at the

same time the seed quality parameters of control seeds stored in cloth bag was declined.

References

- Bradford, K.J. (1986). Manipulation of seeds water relations via osmotic priming to improve germination under stress conditions. Hort. Sci. 59(2): 672-676.
- Doijode, S.D. (1986). Deteriorative changes in okra seeds after artificial ageing Prog. Hort. 18 (3-4): 218-221.
- Delouche, J.C. & Baskin, C.C. (1973). Accelerated aging techniques for predicting relative storability of seed lots. Seed sci. and technol. 1: 427-452.
- Parsley, H.T. (1958). Relation of protoplast permeability of cotton, seed viability and predisposition of seedling diseases. Plt. Disease report. 42:52.
- Karivaratharaju, V., Palaniswamy, V., & Kumarasen, K. (1987). Effect of seed treatment and containers on the storability of brinjal seeds. Seed Res. 129: 141-153.
- McDonald, M. B. (1999). Seed deterioration: physiology, repair and assessment. Seed Sci. and Technol. 27: 177-237.
- Venkatasubramanian, A. & Umarani, R., 2010. Storability of primed seeds of tomato (*Lycopersicon esculentum*), eggplant (*Solanum melongena*) and chili (*Capsicum annum*). Madras Agric. J., 97 (4-6), Pp. 117-124

Received : 26th July 2021

Accepted : 15th October 2022
